

# Asthma Treatment Plan

(This asthma action plan meets NJ Law N.J.S.A. 18A:40-12.8) (Physician's Orders)

**The Pediatric/Adult  
Asthma Coalition  
of New Jersey**  
"Your Pathway to Asthma Control"  
PACNJ approved Plan available at  
www.pacnj.org

Sponsored by  
**AMERICAN  
LUNG  
ASSOCIATION**  
IN NEW JERSEY



**(Please Print)**

|        |                                 |               |                   |
|--------|---------------------------------|---------------|-------------------|
| Name   |                                 | Date of Birth | Effective Date    |
| Doctor | Parent/Guardian (if applicable) |               | Emergency Contact |
| Phone  | Phone                           |               | Phone             |

## HEALTHY



You have **all** of these:

- Breathing is good
- No cough or wheeze
- Sleep through the night
- Can work, exercise, and play

And/or Peak flow above \_\_\_\_\_

**Take daily medicine(s). Some metered dose inhalers may be more effective with a "spacer" - use if directed**

| MEDICINE   | HOW MUCH to take and HOW OFTEN to take it  |
|--|--|
| <input type="checkbox"/> Advair® <input type="checkbox"/> 100, <input type="checkbox"/> 250, <input type="checkbox"/> 500              | _____ 1 inhalation twice a day   |
| <input type="checkbox"/> Advair® HFA <input type="checkbox"/> 45, <input type="checkbox"/> 115, <input type="checkbox"/> 230           | _____ 2 puffs MDI twice a day  |
| <input type="checkbox"/> Alvesco® <input type="checkbox"/> 80, <input type="checkbox"/> 160  | _____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 puffs MDI twice a day   |
| <input type="checkbox"/> Asmanex® Twisthaler® <input type="checkbox"/> 110, <input type="checkbox"/> 220                               | _____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 inhalations <input type="checkbox"/> once or <input type="checkbox"/> twice a day |
| <input type="checkbox"/> Flovent® <input type="checkbox"/> 44, <input type="checkbox"/> 110, <input type="checkbox"/> 220              | _____ 2 puffs MDI twice a day  |
| <input type="checkbox"/> Flovent® Diskus® <input type="checkbox"/> 50 <input type="checkbox"/> 100 <input type="checkbox"/> 250        | _____ 1 inhalation twice a day   |
| <input type="checkbox"/> Pulmicort Flexhaler® <input type="checkbox"/> 90, <input type="checkbox"/> 180                                | _____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 inhalations <input type="checkbox"/> once or <input type="checkbox"/> twice a day |
| <input type="checkbox"/> Pulmicort Respules® <input type="checkbox"/> 0.25, <input type="checkbox"/> 0.5, <input type="checkbox"/> 1.0 | _____ 1 unit nebulized <input type="checkbox"/> once or <input type="checkbox"/> twice a day   |
| <input type="checkbox"/> Qvar® <input type="checkbox"/> 40, <input type="checkbox"/> 80  | _____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 puffs MDI twice a day   |
| <input type="checkbox"/> Singulair® <input type="checkbox"/> 4, <input type="checkbox"/> 5, <input type="checkbox"/> 10 mg             | _____ 1 tablet daily   |
| <input type="checkbox"/> Symbicort® <input type="checkbox"/> 80, <input type="checkbox"/> 160  | _____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 puffs MDI twice a day   |
| <input type="checkbox"/> Other   |  |
| <input type="checkbox"/> None  |  |

Remember to rinse your mouth after taking inhaled medicine.

If exercise triggers your asthma, take this medicine \_\_\_\_\_ minutes before exercise.

## CAUTION



You have **any** of these:

- Exposure to known trigger
- Cough
- Mild wheeze
- Tight chest
- Coughing at night
- Other: \_\_\_\_\_

And/or Peak flow from \_\_\_\_\_ to \_\_\_\_\_

**Continue daily medicine(s) and add fast-acting medicine(s).**

| MEDICINE   | HOW MUCH to take and HOW OFTEN to take it      |
|--|--|
| <input type="checkbox"/> Accunet® <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg                                | _____ 1 unit nebulized every 4 hours as needed |
| <input type="checkbox"/> Albuterol <input type="checkbox"/> 1.25, <input type="checkbox"/> 2.5 mg                                | _____ 1 unit nebulized every 4 hours as needed |
| <input type="checkbox"/> Albuterol <input type="checkbox"/> Pro-Air <input type="checkbox"/> Proventil®                          | _____ 2 puffs MDI every 4 hours as needed      |
| <input type="checkbox"/> Ventolin® <input type="checkbox"/> Maxair <input type="checkbox"/> Xopenex®                             | _____ 2 puffs MDI every 4 hours as needed      |
| <input type="checkbox"/> Xopenex® <input type="checkbox"/> 0.31, <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg | _____ 1 unit nebulized every 4 hours as needed |
| <input type="checkbox"/> Increase the dose of, or add:   |  |
| <input type="checkbox"/> Other   |  |

➡ If fast-acting medicine is needed more than 2 times a week, except before exercise, then call your doctor.

## EMERGENCY



Your asthma is getting worse fast:

- Fast-acting medicine did not help within 15-20 minutes
- Breathing is hard and fast
- Nose opens wide
- Ribs show
- Trouble walking and talking
- Lips blue • Fingernails blue

And/or Peak flow below \_\_\_\_\_

**Take these medicines NOW and call 911. Asthma can be a life-threatening illness. Do not wait!**

|  |   |
|--|---|
| <input type="checkbox"/> Accunet® <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg                                | _____ 1 unit nebulized every 20 minutes |
| <input type="checkbox"/> Albuterol <input type="checkbox"/> 1.25, <input type="checkbox"/> 2.5 mg                                | _____ 1 unit nebulized every 20 minutes |
| <input type="checkbox"/> Albuterol <input type="checkbox"/> Pro-Air <input type="checkbox"/> Proventil®                          | _____ 2 puffs MDI every 20 minutes      |
| <input type="checkbox"/> Ventolin® <input type="checkbox"/> Maxair <input type="checkbox"/> Xopenex®                             | _____ 2 puffs MDI every 20 minutes      |
| <input type="checkbox"/> Xopenex® <input type="checkbox"/> 0.31, <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg | _____ 1 unit nebulized every 20 minutes |
| <input type="checkbox"/> Other   |   |

**Triggers**  
Check all items that trigger patient's asthma:

- Chalk dust
- Cigarette Smoke & second hand smoke
- Colds/Flu
- Dust mites, dust, stuffed animals, carpet
- Exercise
- Mold
- Ozone alert days
- Pests - rodents & cockroaches
- Pets - animal dander
- Plants, flowers, cut grass, pollen
- Strong odors, perfumes, cleaning products, scented products
- Sudden temperature change
- Wood Smoke
- Foods: \_\_\_\_\_
- Other: \_\_\_\_\_

This asthma treatment plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs.

The treatment plan form is a public law, sponsored by the American Lung Association of New Jersey, and the medication is prescribed by a general practitioner. The New Jersey Department of Health, Division of Public Health, has approved this form. The American Lung Association of New Jersey, 400 North 5th Street, Philadelphia, PA 19106, is the national headquarters of the American Lung Association. For more information, contact the American Lung Association at 1-800-558-LUNG.

Although this document has been limited solely to be used by the United States Environmental Protection Agency under Agreement: X9075670-2 to the American Lung Association of New Jersey, it has not been approved through the Agency's public information review process, and therefore, does not necessarily reflect the views of the Agency, and no official endorsement should be inferred.

**REVISED MAY 2009**  
Permission to reproduce blank form  
www.pacnj.org

### FOR MINORS ONLY:

- This student is capable and has been instructed in the proper method of self-administering of the non-nebulized inhaled medications named above in accordance with NJ Law.
- This student is not approved to self-medicate.

Make a copy for patient and for physician file. For children under 18, send original to school nurse or child care provider.

PHYSICIAN/APN/PA SIGNATURE \_\_\_\_\_ DATE \_\_\_\_\_

PARENT/GUARDIAN SIGNATURE \_\_\_\_\_

PHYSICIAN STAMP

## Asthma Treatment Plan Patient/Parent Instructions



The **PACNJ Asthma Treatment Plan** is designed to help everyone understand the steps necessary for the individual patient to achieve the goal of controlled asthma.

### 1. Patients/Parents/Guardians: Before taking this form to your Health Care Provider:

Complete the top left section with:

- Patient's name
- Patient's date of birth
- Patient's doctor's name & phone number
- Parent/Guardian's name & phone number
- An Emergency Contact person's name & phone number

### 2. Your Health Care Provider will:

Complete the following areas:

- The effective date of this plan
- The medicine information for the Healthy, Caution and Emergency sections
- Your Health Care Provider will check the box next to the medication and check how much and how often to take it
- Your Health Care Provider may check "OTHER" and:
  - ❖ Write in asthma medications not listed on the form
  - ❖ Write in additional medications that will control your asthma
  - ❖ Write in generic medications in place of the name brand on the form
- Together you and your Health Care Provider will decide what asthma treatment is best for you or your child to follow.

### 3. Patients/Parents/Guardians & Health Care Providers together:

Discuss and then complete the following areas:

- Patient's peak flow range in the Healthy, Caution and Emergency sections on the left side of the form
- Patient's asthma triggers on the right side of the form
- For Minors Only section at the bottom of the form: Discuss your child's ability to self-administer the inhaled medications, check the appropriate box, and then both you and your Health Care Provider must sign and date the form

### 4. Parents/Guardians: After completing the form with your Health Care Provider:

- Make copies of the Asthma Treatment Plan and give the signed original to your child's school nurse or child care provider
- Keep a copy easily available at home to help manage your child's asthma
- Give copies of the Asthma Treatment Plan to everyone who provides care for your child, for example: babysitters, before/after school program staff, coaches, scout leaders

**This Asthma Treatment Plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs. Not all asthma medications are listed and the generic names are not listed.**

#### Disclaimers:

The use of this Website/PACNJ Asthma Treatment Plan and its content is at your own risk. The content is provided on an "as is" basis. The American Lung Association of the Mid-Atlantic (ALAM-A), the Pediatric/Adult Asthma Coalition of New Jersey and all affiliates disclaim all warranties, express or implied, statutory or otherwise, including but not limited to the implied warranties or merchantability, non-infringement of third parties' rights, and fitness for a particular purpose.

ALAM-A makes no representations or warranties about the accuracy, reliability, completeness, currency, or timeliness of the content. ALAM-A makes no warranty, representation or guaranty that the information will be uninterrupted or error free or that any defects can be corrected.

In no event shall ALAM-A be liable for any damages (including, without limitation, incidental and consequential damages, personal injury/wrongful death, lost profits, or damages resulting from data or business interruption) resulting from the use or inability to use the content of this Asthma Treatment Plan whether based on warranty, contract, tort or any other legal theory, and whether or not ALAM-A is advised of the possibility of such damages. ALAM-A and its affiliates are not liable for any claim, whatsoever, caused by your use or misuse of the Asthma Treatment Plan, nor of this website.

The Pediatric/Adult Asthma Coalition of New Jersey, sponsored by the American Lung Association of New Jersey, and this publication are supported by a grant from the New Jersey Department of Health and Senior Services (NJDHSS), with funds provided by the U.S. Centers for Disease Control and Prevention (USCDCP) under Cooperative Agreement 5U59EH000206-3. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NJDHSS or the USCDCP. Although this document has been funded wholly or in part by the United States Environmental Protection Agency under Agreement XA97256707-2 to the American Lung Association of New Jersey, it has not gone through the Agency's publications review process and therefore, may not necessarily reflect the views of the Agency and no official endorsement should be inferred. Information in this publication is not intended to diagnose health problems or take the place of medical advice. For asthma or any medical condition, seek medical advice from your child's or your health care professional.